


Scientific Note

On the distribution of *Vagrans egista* (Cramer, 1780) in West Bengal, IndiaSobre la distribución de *Vagrans egista* (Cramer, 1780) en el sur de Bengala occidental, IndiaSubhajit Roy^{1*} , Aniruddha Singhamahapatra²  and Suvankar Dutta³

¹Wildlife Information and Nature Guide Society, Durgapur, Paschim Bardhaman, West Bengal 713206, India.  *roy.riju.nitdgp@gmail.com. ²Kadma, Bankura, West Bengal 722151, India. E-mail: aniruddha9735@gmail.com. ³Gopinathpur, Bankura, West Bengal 722101, India. E-mail: duttasuvankar4new@gmail.com

ZooBank: urn:lsid:zoobank.org:pub:C481BD2C-2721-4518-B7E2-B1300161C87B
<https://doi.org/10.35249/rche.47.3.21.09>

Abstract. *Vagrans egista* (Cramer, 1780) is reported for the first time from southern part of West Bengal, India. The Raygar forest of Bankura district in the plateau region is the second locality for the reported species in West Bengal, which extends the distribution range (aerial distance: 525 km) of the species from its known locality, Buxa Tiger Reserve of Alipurduar district in the state. This paper also reports occurrence of *Rapala pheretima* and *Gerosis bhagava* in Bankura district for the first time. Importance of an invasive plant, *Mikania micrantha* has also been discussed as a nectaring plant of butterflies.

Key words: Butterfly; Deccan; invasive plant; nectaring; new record.

Resumen. *Vagrans egista* (Cramer, 1780) es registrada por primera vez en la parte sur de Bengala occidental, India. El bosque de Raygar del distrito de Bankura en la región de la meseta es la segunda localidad reportada en Bengala occidental, lo que extiende el rango de distribución (distancia aérea: 525 km) de la especie desde su localidad conocida, la Reserva de Tigres de Buxa del distrito de Alipurduar en el estado. Esta nota también informa por primera vez la presencia de *Rapala pheretima* y *Gerosis bhagava* en el distrito de Bankura. Se discute la importancia de la planta invasora *Mikania micrantha* como proveedora de néctar para mariposas.

Palabras clave: Deccan; mariposa; néctar; nuevo registro; planta invasora.

India is a megadiverse country in term of ecological diversity, comprising of ten biogeographic zones: Trans Himalaya, Himalaya, Desert, Semi-arid, Western Ghats, Deccan Plateau, Gangetic Plain, Northeast, Islands and Coasts (Rodgers *et al.* 2002) as well as floral and faunal diversity. West Bengal is a state in Eastern India which spans through parts of four –biogeographic zones (Himalaya, Gangetic Plain, Deccan Plateau and Coasts) and is a home to a rich array of biodiversity. Among these biogeographic zones of West Bengal, there is further scope for exploration in the plateau region. The present study site (Fig. 1) is Raygar forest near Sutan (22°50'02.7" N 86°43'19.0" E; altitude: 214 m) at Ranibandh block in undulating south-western part of Bankura district of Medinipur division. The area is covered with a dense dry deciduous forest (Mukherjee and Mondal 2020), primarily dominated by some common trees as *Shorea robusta* Roth, *Diospyros melanoxylon* Roxb., *Madhuca longifolia* (J.Konig) J.F.Macbr., *Dendrocalamus strictus* (Roxb.) Nees, *Butea monosperma*

Received 23 May 2021 / Accepted 16 July 2021 / Published online 31 August 2021
Responsible Editor: José Mondaca E.

(Lam.) Taub., *Bombax ceiba* L., *Phoenix sylvestris* (L.) Roxb., *Ficus* spp. of deciduous forest, which is bisected by a rain-fed seasonal stream. The region is characterized by hard beds of primary laterite formed by in situ weathering of Gneissic rocks (Sen *et al.* 2020) and the ferruginous laterite crust overlain with latosol is rich in sandy loam (Ghosh and Guchhait 2015). Such typical lateritic type soil of Rarh region holds the last remnants of dry deciduous forests (Ghosh 2008). This part of Bankura district is connected with Chhotanagpur Plateau (the north-easternmost extension of Deccan Plateau) in the West and the lower gangetic plains in the East. This rolling upland is an eroded remnant of the Chhotanagpur Plateau. Tropical dry sub-humid climatic conditions prevail in the study area, which is classified as Zone Aw according to Köppen-Geiger climatic classification (Peel *et al.* 2007). The average temperature is 25.7 °C: May and January represent the hottest (41 °C) and coolest (6 °C) months respectively; and the average annual rainfall is 1455 mm: July and January are the wettest (505.7 mm) and driest months (7.2 mm) respectively (Anonymous 2021).



Figure 1. Study site from Bankura district, West Bengal, India (Source: Google Earth maps). / Sitio de estudio en el distrito de Bankura, Bengala Occidental, India (Fuente: mapas de Google Earth).

Butterflies are good indicators of ecosystem health due to their sensitivity to ecological parameters (New 1991; Bonebrake *et al.* 2010). They are one of the most important pollinators and herbivores in nature (Kunte 2000). The abundance of terrestrial insects including butterflies are declining by 8.81% in each decade, even more in the unprotected areas (van Klink *et al.* 2020). Hence, the study on the butterflies is significant to maintain ecological balance. The authors are carrying extensive field surveys throughout heterogeneous patches of Bankura district since 2015 as part of a long-term study on the butterfly diversity of plateau region. During this study, an individual of *Vagrans egista* (Cramer, 1780) has been observed to nectaring on *Mikania micrantha* Kunth flower-heads and the butterfly has also been photographed. The butterfly was identified comparing the captured photographs with standard literatures (Evans 1932; Kehimkar 2016).

Observation. An individual of *Vagrans egista* has been observed to nectaring on flower of *Mikania micrantha* at about 01:26 PM (GMT: +5:30) on 15.ii.2020 (Figs. 2-3). Two of the authors (SD and AS) captured the photographs of the butterfly and identified the species comparing the photos with aforementioned literatures. It is a distinctive species with its golden tawny upper side with its narrow dark brown outer edge (Kehimkar 2016) and the underside has pale purple shimmering with a series of small glowing white spots (Cramer 1780). Besides, *Phalanta phalantha* (Drury, 1773) were also observed to nectaring on *Mikania micrantha* flowers in association with *Vagrans egista*. In addition, *Rapala pheretima* (Hewitson, 1863) and *Gerosis bhagava* (Moore, 1865) (Figs. 4-5) are also reported first time from the Bankura district. Therefore, 25 more species has been observed nectaring on the flowers of *Mikania micrantha* (Table 1).



Figures 2-5. Butterflies nectaring on *Mikania micrantha*. 2-3. *Vagrans egista*, Raygar forest, India. 4-5. *Rapala pheretima* and *Gerosis bhagava*, newly recorded species for Bankura district. / 2-3. *Vagrans egista* libando néctar sobre *Mikania micrantha*, bosque de Raygar, India. / 4-5. *Rapala pheretima* y *Gerosis bhagava*, nuevos registros de especies para el distrito de Bankura.

Table 1. Systematic list of observed nectaring Lepidoptera on flowers of *Mikania micrantha*. / Lista sistemática de los lepidópteros nectarizantes observados en flores de *Mikania micrantha*.

Sl. No.	Species	No. of Individuals
Family Scythrididae		
1	<i>Eretmocera impactella</i> Walker, 1864	5
Family Hesperidae		
2	<i>Ampittia dioscorides</i> (Fabricius, 1793)	1
3	<i>Gerosis bhagava</i> (Moore, 1866)	1
4	<i>Pelopidas mathias</i> (Fabricius, 1798)	1
5	<i>Parnara bada</i> (Moore, 1878)	3
Family Pieridae		
5	<i>Catopsilia pyranthe</i> (Linnaeus, 1758)	6
6	<i>Delias eucharis</i> (Drury, 1773)	3
7	<i>Eurema blanda</i> (Boisduval, 1836)	1
8	<i>Eurema hecabe</i> (Linnaeus, 1758)	2
Family Lycaenidae		
9	<i>Spindasis ictis</i> (Hewitson, 1865)	1
10	<i>Spindasis vulcanus</i> (Fabricius, 1775)	6
11	<i>Amblypodia anita</i> Hewitson, 1862	4
12	<i>Tajuria cippus</i> (Fabricius, 1798)	1
13	<i>Rapala manea</i> (Hewitson, 1863)	3
14	<i>Rapala pheretima</i> (Hewitson, 1863)	6
15	<i>Rapala varuna</i> (Horsfield, 1829)	3
16	<i>Anthene emolus</i> (Godart, 1824)	2
Family Nymphalidae		
17	<i>Euploea core</i> (Cramer, 1780)	5
18	<i>Junonia almana</i> (Linnaeus, 1758)	6
19	<i>Junonia altites</i> (Linnaeus, 1763)	3
20	<i>Junonia iphita</i> (Cramer, 1779)	2
21	<i>Tanaecia lepidea</i> (Butler, 1868)	85
22	<i>Athyma perius</i> (Linnaeus, 1758)	2
23	<i>Neptis jumbah</i> Moore, 1858	2
24	<i>Pantoporia hordonia</i> (Stoll, 1790)	4
25	<i>Phalanta phalantha</i> (Drury, 1773)	12
26	<i>Vagrans egista</i> (Cramer, 1780)	1
Family Noctuidae		
27	<i>Episteme adulatrix</i> (Kollar, 1844)	1
Family Erebidae		
28	<i>Mangina argus</i> (Kollar, 1844)	1

Before this report, *Vagrans egista* were reported only from Buxa Tiger Reserve (BTR) of Alipurduar district in West Bengal (Sinha *et al.* 2019). Hence, this record is the first report from southern part of the state and second locality report from West Bengal. The present locality is around 500 km (aerial distance) away from previously reported nearby locality at BTR of Alipurduar district in north-eastern part of West Bengal. Moreover, the species is distributed from Uttarakhand to Northeast India and West Bengal, Odisha, Jharkhand, Himachal Pradesh, Andhra Pradesh and Chhattisgarh in India (van Gasse 2013; Varshney and Smetacek 2015; Kehimkar 2016; Kirti *et al.* 2016; Payra *et al.* 2016; Sisodia and Naidu 2019; Gokhale 2021). Globally, *Vagrans egista* is distributed in Nepal, India, Bangladesh, south China to South-east Asia, *i.e.*, Myanmar, Thailand, Laos, Cambodia, Vietnam, Philippines and parts of Indonesia and Malaysia (Kehimkar 2016; Inayoshi 2021). This observation fills the missing sampling gap of the species from the plateau part of West Bengal. Since this is not known to be a migratory species, a resident population of the species can be assumed in this region. Besides, literature search reveal that report of *Rapala pheretima* and *Gerosis bhagava* are the first records from Bankura district (Nayak 2020; Mukherjee and Mondal 2020).

The *Mikania micrantha* were also observed as notably significant plant for butterfly fauna during the study. This is an extremely fast-growing climber native to Central and South America (Wirjahar 1976). It is now distributed worldwide and is considered as one of the world's most notorious invaders (Holm *et al.* 1977), introduced in Indian subcontinent by the British in 1940s to camouflage airfields during World War II (Tripathi *et al.* 2011) and as ground cover in tea plantations (Prabu *et al.* 2014). Flowering season of the plant in eastern and north-east India is from October to late November and these annual plants die in December and January (Swamy and Ramakrishnan 1987). This plant is known to cause disastrous ecological effects as it kills nearby plant species by reducing light beneath its canopy (Huang *et al.* 2000). However, while many species suffer from the invasion of a species, some of them also benefit from it (Davis *et al.* 2018). Likely, *Mikania micrantha* is an especially important nectaring plant for the flower-visiting insects such as butterflies, moths, hoverflies, bees, and wasps (Hong *et al.* 2011). Thus, this climber enhances the abundance and diversity of insects during the period of its bloom. To control growth of such invasive but beneficiary plant to the insects, biological control might be adopted during blooming season rather than use of herbicide and forest fire. For instance, use of *Ipomoea batatas* (L.) or *Cuscuta campestris* Yunck might be attempted for biological control of *Mikania micrantha* (Shen *et al.* 2015; Li *et al.* 2012). Monoculture of exotic species during afforestation should be avoided, which is responsible for decline of diversity and abundance of flower-visiting insects including butterflies (Nagulapalli *et al.* 2020). However, further extensive study should be attempted to protect butterflies as well as their host and nectaring plants of the plateau region of West Bengal.

Acknowledgement

The authors acknowledge Mr. Amar Kumar Nayak and Mr. Rajib Dey for their constant encouragement in preparing the manuscript. Dr. Roger Kendrick, Dr. Shantanu Jha, Mr. Balakrishnan Valappil and Dr. Priyantha Wijesinghe are heartily acknowledged for discussions which have enriched the knowledge of the authors about nectaring. The authors also express their gratitude to their family members for constant motivation for the field surveys.

Literature Cited

Anonymous (2021) Monthly Climate Averages in Ranibandh, West Bengal, India. World Weather Online. Accessed date May 21, 2021. Available at: <https://www.worldweatheronline.com/ranibandh-weather-averages/west-bengal/in.aspx>

- Bonebrake, T.C., Ponisio, L.C., Boggs, C.L. and Ehrlich, P.R. (2010)** More than just indicators: A review of tropical butterfly ecology and conservation. *Biological Conservation*, 143(8): 1831-1841. <https://doi.org/10.1016/j.biocon.2010.04.04>
- Cramer, P. (1780)** De uitlandsche kapellen, voorkomende in de drie waereld-deelen Asia, Africa en America (Papillons exotiques des trois parties du monde L'Asie, L'Afrique et L'Amerique). Concordia et Libertate, Amsterdam, The Netherlands, 3 (23-24): 158 pp; pl. 281, f. C, D.
- Davis, E.S., Kelly, R., Maggs, C.A. and Stout, J.C. (2018)** Constrasting impacts of highly invasive plant-species on flower-visiting insect communities. *Biodiversity and Conservation*, 27(8): 2069-2085. <https://doi.org/10.1007/s10531-018-1525-y>
- Evans, W.H. (1932)** The Identification of Indian Butterflies. Bombay Natural History Society, Mumbai, India. 454 pp.
- Ghosh, A. (2008)** Ethnomedicinal plants used in West Rarrh region of West Bengal. *Indian Journal of Natural Products and Resources*, 7(5): 461-465.
- Ghosh, S. and Guchhait, S.K. (2015)** Characterization and evolution of primary and secondary laterites in northwestern Bengal basin, West Bengal, India. *Journal of Palaeogeography*, 4(2):203-230. <https://doi.org/10.3724/SP.J.1261.2015.00074>
- Gokhale, P. (2021)** *Vagrans egista* (Cramer, [1780]) - Vagrant. Kunte, K., Sondhi, S. and Roy, P. (Chief Editors), *Butterflies of India*, v. 3.11. Indian Foundation for Butterflies. Accessed May 17, 2021. Available at: <http://www.ifoundbutterflies.org/sp/908/Vagrans-egista>
- Huang, Z.L., Cao, H.L., Liang, X.D., Ye, W.H., Feng, H.L. and Cai, C.X. (2000)** The growth of damaging effects by *Mikania micrantha* in different habitats. *Journal of Tropical and Subtropical Botany*, 8: 131-138.
- Holm, L.G., Plucknett, D.L., Pancho, J.V. and Herberger, J.P. (1977)** The World's worst weeds. Distribution and Biology. University Press of Hawaii, Honolulu, United States of America, 610 pp.
- Hong, L., Shen, H., Ye, W.H. and Cao, H.L. (2011)** Study of pollinating insects of *Mikania micrantha* H.B.K and their foraging behavior. *Journal of South China Normal University (Natural Science Edition)*, 43: 98-102. <https://doi.org/10.6054/j.jscnun.2011.11.016>
- Inayoshi, Y. (2021)** A Check List of Butterflies in Indo – China (chiefly from Thailand, Laos & Vietnam). Accessed May 21, 2021. Available at: <http://yutaka.it-n.jp/hel/710140010.html>
- Kehimkar, I. (2016)** Butterflies of India. Bombay Natural History Society, Mumbai, India, 258 pp.
- Kirti, J.S., Mehra, D. and Sidhu, A.K. (2016)** *Vagrans egista sinha* (Kollar, 1844) (Lepidoptera: Papilionoidea: Nymphalidae): range extension and an addition to the butterfly fauna of Himachal Pradesh, India. *Check List*, 12(4)b: 1923. <https://doi.org/10.15560/12.4.1923>
- Kunte, K. (2000)** India - A Landscape: Butterflies of Peninsular India. Universities Press, Hyderabad, India, 147 pp.
- Li, F.L., Li, M.G., Zan, Q.J., Qiang, G.U.O., Zhang, W.Y., Zhi, W.U. and Wang, Y.J. (2012)** Effects of the residues of *Cuscuta campestris* and *Mikania micrantha* on subsequent plant germination and early growth. *Journal of Integrative Agriculture*, 11(11): 1852-1860.
- Mukherjee, K. and Mondal, A. (2020)** Butterfly diversity in heterogeneous habitat of Bankura, West Bengal, India. *Journal of Threatened Taxa*, 12(8): 15804-15816. <https://doi.org/10.11609/jott.5136.12.8.15804-15816>
- Nagulapalli, S.L., Kunal, G., Das, R. and Jha, S. (2020)** Insect Pollinators and their abundance in West Bengal. *Indian Journal of Entomology*, 82(4): 849-853. <https://doi.org/10.5958/0974-8172.2020.00110.8>
- Nayak, A. (2020)** A checklist of butterfly fauna from Bankura town, West Bengal, India. *Journal of Threatened Taxa*, 12(13): 16868-16878. <https://doi.org/10.11609/jott.4882.12.13.16868-16878>

- New, T.R. (1991) Butterfly Conservation. Oxford University Press. Oxford, United Kingdom, 224 pp.
- Payra, A., Das, G.N., Boruah, B., Dash, S.K., Das, U.P. and Sethy, J. (2016) Butterfly diversity in two selected fringe area of Simlipal Biosphere Reserve, Odisha, India, with notes on some important sightings. *Journal of Wildlife Research*, 4(2): 17-25.
- Peel, M.C., Finlayson, B.L. and McMahon, T.A. (2007) Updated world map of the Köppen-Geiger climate classification. *Hydrology and Earth System Sciences*, 11: 1633-1644. <https://doi.org/10.5194/hess-11-1633-2007>
- Prabu, N.R., Stalin, N. and Swamy, P.S. (2014) Ecophysiological attributes of *Mikania micrantha*, an exotic invasive weed, at two different elevations in the tropical forest regions of the Western Ghats, South India. *Weed Biology and Management*, 14(1): 59-67.
- Rodgers, W.A., Panwar, H.S. and Mathur, V.B. (2002) Wildlife Protected Areas in India: A review (Executive Summary), Wildlife Institute of India, Dehradun, India, 51 pp.
- Sen, S., Chakraborty, D., Chakraborty, J. and Das, K. (2020) Application of Geospatial Techniques for Land Resource Characterization and Management of Micro-watersheds: A Study in the Lateritic Micro-watersheds of River Kangshabati, Block Ranibandh, District Bankura. Recent Trends in Agriculture, Environment, Sustainability and Life Sciences. pp. 126-155, Sparrow Publication, Kolkata, India.
- Shen, S., Xu, G., Clements, D.R., Jin, G., Chen, A., Zhang, F. and Kato-Noguchi, H. (2015) Suppression of the invasive plant mile-a-minute (*Mikania micrantha*) by local crop sweet potato (*Ipomoea batatas*) by means of higher growth rate and competition for soil nutrients. *BMC Ecology*, 15(1): 1.
- Sinha, R.K., Sengupta, S., Roy, A.B., Baidya, S., Roy, S., Das, N., Paul, A. and Pal, A. (2019) Butterflies of Buxa Tiger Reserve, Citadel, Kolkata, India, 248 pp.
- Sisodia, A. and Naidu, R. (2019) Extension of the known distribution of the Vagrant butterfly, *Vagrans egista* (Cramer, [1780]) (Lepidoptera: Nymphalidae) to Bastar, Chhattisgarh. *Bionotes*, 21(1): 12-13.
- Swamy, P.S. and Ramakrishnan, P.S. (1987) Effect of fire on population dynamics of *Mikania micrantha* H.B.K. during early succession after slash and burn agriculture (jhum) in North-east India. *Weed Research*, 27: 397-403.
- Tripathi, R.S., Khan, M.L. and Yadav, A.S. (2011) Biology of *Mikania micrantha* H.B.K.: a review. Invasive Alien Plants: an ecological appraisal for the Indian subcontinent. (ed. Bhatt, J.R., Singh, J.S., Singh, S.P., Tripathi, R.S. and Kohli, R.K.), pp. 99-107, Centre for Agriculture and Bioscience International, Willingford, United Kingdom.
- van Gasse, P. (2013) Butterflies of India - Annotated Checklist. Accessed May 22, 2021. Available at: https://www.ifoundbutterflies.org/images/PaulVanGasse/Butterflies_of%20India-Annotated_checklist-1.pdf
- van Klink, R., Bowler, D.E., Gongalsky, K.B., Swengel, A.B., Gentile, A. and Chase, J.M. (2020) Meta-analysis reveals declines in terrestrial but increases in freshwater insect abundances. *Science*, 368(6489): 417-420. <https://doi.org/10.1126/SCIENCE.AAX9931>
- Varshney, R.K. and Smetacek, P. (2015) A synoptic catalogue of the Butterflies of India, Butterfly Research Centre, Bhimtal, India and Indinov Publishing, New Delhi, India, 261 pp. <https://doi.org/10.13140/RG.2.1.3966.2164>
- Wirjajhar, S. (1976) Autoecological study of *Mikania* spp. Proceedings of the Fifth Asian Pacific Weed Science Society Conference (ed. Kataoko, T., Furuya, S., Hyakutake, H., Ishizuka, K., Matsunaka, S. and Takabayashi, M.), Tokyo, Japan, pp. 70-73.