

Scientific Note

New record of *Mythapion adesmiae* Kissinger, 2005 (Coleoptera: Brentidae) in Argentina

Nuevo registro de *Mythapion adesmiae* Kissinger, 2005 (Coleoptera: Brentidae) en Argentina

Sabrina S. Gavini^{1*}  and Marcelo E. Kun² 

¹INIBIOMA, CONICET-Centro Regional Universitario Bariloche, Universidad Nacional del Comahue, Quintral 1250, Bariloche, Río Negro, Argentina.  *sabrinagavini@comahue-conicet.gob.ar. ²Laboratorio de Zoología, Centro Regional Universitario Bariloche, Universidad Nacional del Comahue, Quintral 1250, Bariloche, Río Negro, Argentina. E-mail: marcelo.kun@crub.uncoma.edu.ar

ZooBank: urn:lsid:zoobank.org:pub:B13EDF04-2CB3-4A06-B9D8-C652E2D561AE
<https://doi.org/10.35249/rche.48.4.22.06>

Abstract. The weevil *Mythapion adesmiae*, previously known from Chilean Patagonia (Aysén and Magallanes regions) and the province of Santa Cruz in Argentina, is recorded for the first time for the province of Río Negro, above the treeline, between 1744-1850 m a.s.l. on Cerro Challhuaco associated with *Adesmia corymbosa*, which is a new plant association. The known records suggest an extensive distribution along the Patagonian Andes of Chile and Argentina.

Key words: Apioninae; Fabaceae; Patagonia; vegetación alto-andina.

Resumen. El gorgojo *Mythapion adesmiae* conocido previamente para la Patagonia chilena (regiones de Aysén y Magallanes) y la provincia de Santa Cruz en Argentina, es registrado por primera vez para la provincia de Río Negro, por encima de los bosques Andino-Patagónicos, a 1744-1850 m en el Cerro Challhuaco, asociado con plantas de *Adesmia corymbosa*, representando una nueva asociación. Los registros conocidos sugieren una amplia distribución a lo largo de los Andes Patagónicos de Chile y Argentina.

Palabras clave: Apioninae; Fabaceae; high-Andean vegetation; Patagonia.

Weevils are beetles belonging to Curculionoidea, the largest superfamily of Coleoptera with more than 70,000 species in the world fauna (Oberprieler *et al.* 2007; Bouchard *et al.* 2009). Excepting Scolytinae, Curculionoidea species are usually recognized by their modified head into a rostrum. The subfamily Apioninae (Brentidae *sensu latu*, straight-snouted weevils) is a monophyletic group (Wanat 2001; Marvaldi *et al.* 2002; Alonso-Zarazaga and Wanat 2014) of very small (0.75-13.0 mm in length) pear-shaped weevils, with straight antennae, having eight narrow segments and a widened compact club, and elongate cylindrical trochanters (Kissinger 2005). Both larvae and adults feed on living plant tissues (stems, leaves, roots, inflorescences) as well as fruits and seeds. In addition, larvae often develop inside stems or roots (forming gall-like structures), leaves, and even seeds (Marvaldi and Lanteri 2005). Even though some Apioninae are associated with gymnosperms (Kissinger 1968; Alonso-Zarazaga and Wanat 2014), most lineages

Received 5 October 2022 / Accepted 20 October 2022 / Published online 28 October 2022

Responsible Editor: José Mondaca E.



Este es un artículo de acceso abierto distribuido bajo los términos de la licencia Creative Commons License (CC BY NC 4.0)

are associated with angiosperms of the families Apiaceae, Asteraceae, Euphorbiaceae, Fabaceae and Lamiaceae, among other families (Anderson and Kissinger 2002; Badenes-Perez and Johnson 2007; Maia 2012; Alonso-Zarazaga and Wanat 2014). The evolutionary success of these weevils apparently lies in their specialized endophytophagy (Brooks and McLennan 2002; Oberprieler *et al.* 2007).

The knowledge of Apioninae and their distribution in South America is limited; nonetheless, we can highlight the studies in Chile (Kissinger 2002, 2005; Elgueta and Marvaldi 2006) and a recent review of the taxonomic diversity of Brazilian Apioninae (de Sousa *et al.* 2019). The genus *Mythapion* Kissinger, 2005 encompass few species with scarce biological and ecological information. The species recorded for Argentina are *M. trifolianum* Kissinger, *M. simplex* Béguin-Billecocq (*Apion simplex*), *M. rufonigrum* Béguin-Billecocq (*Apion rufonigrum*) (Kissinger 2005; de Sousa *et al.* 2019), and *Mythapion adesmiae* Kissinger, 2005, recorded for Santa Cruz province and in Chile (Kissinger 2005; Elgueta and Marvaldi 2006). Herein we provide a new record for Río Negro province.




Figure 1. *Mythapion adesmiae* Kissinger records in Argentina (A). Santa Cruz province in 1998, in proximity to the Torres del Paine National Park, and Río Negro province in 2019 in the Nahuel Huapi National Park. The Andean environment above treeline where specimens were collected (B, C). Host plant *Adesmia corymbosa* (D). Credit of *Adesmia* photograph, Marcela Ferreyra, used with permission. Photographs of non-damaged seeds (left side) and damaged seeds (right side) (E). Seeds are about 3 mm in diameter. / Registros de *Mythapion adesmiae* Kissinger en Argentina (A). Provincia de Santa Cruz en 1998, en cercanías al Parque Nacional Torres del Paine, y provincia de Río Negro en 2019 en el Parque Nacional Nahuel Huapi. El ambiente andino por encima del bosque donde se recolectaron los ejemplares (B, C). Planta huésped *Adesmia corymbosa* (D). Crédito de la fotografía de *Adesmia* a Marcela Ferreyra, usada con autorización. Fotografías de semillas no dañadas (lado izquierdo) y semillas dañadas (lado derecho) (E). Las semillas tienen unos 3 mm de diámetro.

According to web site Coleoptera-neotropical.org the first and unpublished detection of *M. adesmiae* in the Argentine Patagonia occurred on December of 1998 in the region of Torres del Paine, located in the transition area between the Magellanic forests and the Patagonian steppes (collected by Juan E. Barriga-Tuñón from Universidad Católica del Maule, Chile). Those specimens were found in Chile and Santa Cruz province, Argentina (Fig. 1A). In February of 2019 we found specimens of *M. adesmiae* on Cerro Challhuaco, between 1744-1850 m a.s.l. ($41^{\circ}16'S$ $71^{\circ}18'W$), in the vicinity of San Carlos de Bariloche within the Nahuel Huapi National Park, Río Negro province (Fig. 1A). Deciduous forests of *N. pumilio* occur from ~1000 m to the treeline that is about 1600 m, where this tree grows prostrated and forms a low 'krummholz'. Above the treeline unfolds the high-Andean ecosystem characterized by a sparse and low vegetation cover (< 20% at 1600 m vs. < 5% at 2000 m, see Figs. 1B, C) resulting in a patchy-landscape represented by isolated cushion-plant patches and a surrounding bare ground matrix (Nuñez *et al.* 1999; Gavini *et al.* 2020). Mean annual temperature is $3.7^{\circ}C$, with short, mild, dry summers and cold, wet winters, whereas mean annual precipitation is approx. 860 mm, mostly in the form of snow between May and October (Gavini *et al.* 2020). Asteraceae, Poaceae and Apiaceae are the best represented families, with perennial herbs, dwarf shrubs and cushion nursing plants the predominant life forms (Ezcurra and Gavini 2020). Individuals of *M. adesmiae* (Fig. 2) were found associated with *Adesmia corymbosa* Clos (Fabaceae) (Fig. 1D), mainly within fruits and newly opened flowers (2 specimens), with their bodies covered with pollen. Additionally, seeds showed signs of predation (Fig. 1E). The specimens' identification was done following Kissinger (2005) and the voucher specimens (24 individuals) were deposited in the Zoology Department at the Centro Regional Universitario Bariloche, Río Negro, Argentina.




Figure 2. *Mythapion adesmiae* Kissinger. Dorsal view (A). Lateral view (B). Length of adult female specimens was 1.93 mm - 1.99 mm (n=3) (length of the rostrum in females: 0.60 mm - 0.62 mm), length of adult male specimens was 1.89 mm - 2.04 mm (n=5) (length of the rostrum in males: 0.53 mm - 0.55 mm). Scale: 500 μ m. / *Mythapion adesmiae* Kissinger. Vista dorsal (A). Vista lateral (B). La longitud de hembras adultas fue de 1,93 mm - 1,99 mm (n=3) (longitud del rostro en hembras: 0,60 mm - 0,62 mm), la longitud de machos adultos fue de 1,89 mm - 2,04 mm (n=5) (longitud del rostro en machos: 0,53 mm - 0,55 mm). Escala: 500 μ m.

The known records suggest that *M. adesmiae* is probably distributed throughout the entire Patagonian Andes, possibly reaching Neuquén province. Furthermore, the species survive the extreme climatic conditions of cold, freeze and drought that characterize the high-Andean environment (Chown 1993; Van der Merwe *et al.* 1997). *Mythapion adesmiae* is associated with the genus *Adesmia* (Fabaceae), such as *A. boronioides* and *A. pegajosa* (Kissinger 2005) being *A. corymbosa* a new record. There are about 100 species of *Adesmia* in Argentina, herbs and shrubs that grow in montane and semi-desert zones (Ulibarri and Burkart 2000), such as *A. glomerula*, *A. longipes* and *A. parvifolia* from the high Andes of Patagonia (Ferreyra *et al.* 2020) that could be also host plants of *M. adesmiae*. Despite their seed-consuming behavior, these weevils may be pollinators because when they reach inflorescences for feeding, shelter, mating, and oviposition they transport pollen and pollinate flowers (Eriksson 1994). Pollination by weevils could be important in environments characterized by low abundance and restricted activity of flying pollinating insects (Körner 2003).

Acknowledgments

This study was financed by grants to Marcelo Kun, Project 04/215 - Arthropod diversity in Argentine Patagonia and their relationships with animals, plants and man - provided by the Universidad Nacional del Comahue, CRUB. We are grateful to the Administración de Parques Nacionales de Argentina that provided permission to work at Cerro Challhuaco.

Literature Cited

- Alonso-Zarazaga, M.A. and Wanat, M. (2014)** Apioninae Schoenherr, 1823. In: Leschen, R.A.B., and Beutel, R.G. (Eds.), *Coleoptera, Beetles. Vol. 3. morphology and systematics (Phytophaga)*. In: Kristensen, N.P. and Beutel, R.G. (Eds.), *Handbook of zoology. Arthropoda: Insecta*. Walter de Gruyter, Berlin, Boston, 395-415 pp.
- Anderson, R.S. and Kissinger, D.G. (2002)** Brentidae Billberg 1820. In: Arnett, R.H Jr., Thomas, M.C, Skelley, P.E., and Frank, J.H. (Eds.), *American Beetles Vol. 2. Polyphaga: Scarabaeoidea through Curculionoidea*. CRC Press, Boca Raton, 711-719 pp.
- Badenes-Perez, F.R. and Johnson, M.T. (2007)** Ecology and impact of *Allorhogas* sp. (Hymenoptera: Braconidae) and *Apion* sp. (Coleoptera: Curculionoidea) on fruits of *Miconia calvescens* DC (Melastomataceae) in Brazil. *Biological Control*, 43: 317-322.
- Bouchard, P., Grebennikov, V.V., Smith, A.B.T. and Douglas, H. (2009)** *Biodiversity of Coleoptera*. Insect Biodiversity. Science and Society. Chichester, UK: Blackwell Publishing Ltd., 265-301 pp.
- Brooks, D.R. and McLennan, D.A. (2002)** *The Nature of Diversity: An Evolutionary Voyage of Discovery*. University of Chicago Press, Chicago, 676 pp.
- Chown, S.L. (1993)** Desiccation resistance in six sub-antarctic weevils (Coleoptera: Curculionidae): Humidity as an abiotic factor influencing assemblage structure. *Functional Ecology*, 7(3): 318-325.
- Coleoptera neotropical**. Available at: <http://coleoptera-neotropical.org>. Accessed: October 2022.
- De Sousa, W.O., Ribeiro-Costa, C.S. and Rosado-Neto, G.H. (2019)** A preliminary overview of the Brazilian Apioninae (Coleoptera: Brentidae) with an illustrated key for genera, and a checklist with distribution information. *Biota Neotropica*, 19(4): e20190813.
- Elgueta, M. and Marvaldi, A.E. (2006)** Lista sistemática de las especies de Curculionoidea (Insecta: Coleoptera) presentes en Chile, con su sinonimia. *Boletín del Museo Nacional de Historia Natural, Chile*, 55: 113-153.
- Eriksson, R. (1994)** The remarkable weevil pollination of the neotropical Carludovicoideae (Cyclanthaceae). *Plant Systematics and Evolution*, 189: 75-81.

- Ezcurra, C. and Gavini, S.S. (2020)** Alpine Plant Diversity in Temperate Mountains of South America. Encyclopedia of the World's Biomes. In: Goldstein, M.I., DellaSala, D.A. (Eds.). Pp. 323-334. Elsevier.
- Ferreyyra, M., Ezcurra, C. and Clayton, S. (2020)** *Flores de Alta Montaña de los Andes Patagónicos / High Mountain Flowers of the Patagonian Andes*. Second Edition. La Plata, Argentina. 72 pp.
- Gavini, S.S., Ezcurra, C. and Aizen, M.A. (2020)** Patch-level facilitation fosters high-Andean plant diversity at regional scales. *Journal of Vegetation Science*, 31: 1135-1145.
- Kissinger, D.G. (1968)** *Curculionidae subfamily Apioninae of North and Central America with reviews of the world genera of Apion Herbst (Coleoptera)*. Taxonomic Publications. South Lancaster, Massachusetts. 559 pp.
- Kissinger, D.G. (2002)** A new genus of Apionidae from Chile and Argentina. *The Coleopterists Bulletin*, 56(2): 315-316.
- Kissinger, D.G. (2005)** Review of Apioninae of Chile (Coleoptera: Curculionoidea: Apionidae). *The Coleopterists Bulletin*, 59(1): 71-90.
- Körner, C. (2003)** *Alpine plant life: functional plant ecology of high mountain ecosystems*. 2nd ed. Heidelberg: Springer. Berlin, Germany. 500 pp.
- Maia, C.M. (2012)** Coleopterous galls from the Neotropical Region. *Papéis Avulsos de Zoologia*, 52(15): 175-184.
- Marvaldi, A.E. and Lanteri, A.A. (2005)** Key to higher taxa of South American weevils based on adult characters (Coleoptera, Curculionoidea). *Revista Chilena de Historia Natural*, 78: 65-87.
- Marvaldi, A.E., Sequeira, A.S., O'Brien, C.W. and Farrell, B.D. (2002)** Molecular and morphological phylogenetics of weevils (Coleoptera, Curculionoidea): Do niche shifts accompany diversification? *Systematic Biology*, 51(5): 761-785.
- Nuñez, C.I., Aizen, M.A. and Ezcurra, C. (1999)** Species associations and nurse plant effects in patches of high-Andean vegetation. *Journal of Vegetation Science*, 10: 357-364.
- Oberprieler, R.G., Marvaldi, A.E. and Anderson, R.S. (2007)** Weevils, weevils, weevils everywhere. *Zootaxa*, 1668: 491-520.
- Ulibarri, E.A. and Burkart, A. (2000)** Sinopsis de las especies de *Adesmia* (Leguminosae, Adesmiaeae) de la Argentina. *Darwiniana*, 38: 59-126.
- Van der Merwe, M., Chown, S. and Smith, V. (1997)** Thermal tolerance limits in six weevil species (Coleoptera, Curculionidae) from sub-Antarctic Marion Island. *Polar Biology*, 18: 331-336.
- Wanat, M. (2001)** *Genera of Australo-Pacific Rhadinocybinae and Myrmacelinae with biogeography of the Apionidae (Coleoptera: Curculionoidea) and phylogeny of the Brentidae (s. lato)*. Mantis, Olsztyn. 432 pp.